

CONTROL OF BED BUGS

Students suspected of carrying bed bugs (*Cimex lectularius*) will be discretely removed from the classroom so that the school nurse or another qualified individual can perform an inspection of the student's clothing and belongings, including shoes, jackets, hats, books, backpacks, school supplies, etc.

After a positive identification, the school administration will directly contact the student's parent or guardian and provide the following information:

- A. Instruct the parent or guardian to send the student to school with a sealable plastic bag containing a clean, freshly laundered change of clothes. The clothes should be washed at the hottest recommended setting and tumble dried on high heat for at least 30 minutes before being placed in the bag.
- B. Request that the parent or guardian send only essential items to school with the student. These items should be inspected upon arrival at school. The school may offer to keep non-essential items overnight to help ensure the items are bed bug free.
- C. Suggest keeping school items sealed in a plastic bag at home and limit items going back and forth from home to school until the infestation is under control.

The Administration will also send to the parents or guardians of all students in the classroom of the student identified as carrying bed bugs a notification letter and a bed bug "Fact Sheet," which will provide steps parents or guardians should take to address a bed bug infestation in their place of residence.

To help minimize the potential transfer of bed bugs or their eggs from one student's belongings to another, students identified as carrying bed bugs will have their personal belongings (coat, backpack, lunch bag, etc.) separated from those of their classmates or placed in individual plastic containers labeled with the students' names.

A school nurse will be assigned to recheck students previously identified as carrying bed bugs on the morning the students return to school. If bed bugs are found, the Administration will contact the student's parents or guardians, as well as the Cuyahoga County Board of Health, or any other appropriate public agency, as deemed necessary by the school principal.

Because bed bugs are not known to transmit disease, students identified as potentially carrying bed bugs will not be excluded from school. However, all reasonable steps to safely remove bed bugs from such students and prevent the spread of future infestations may be taken. The Administration will discretely follow the guidelines provided in this policy with the affected student's dignity in mind.

Adopted: August 28, 2017

School Letterhead

Date

Dear Parent or Guardian:

This letter has been sent to you in response to a bed bug being found in your child's classroom. Although it is often very difficult to identify the source of the bed bug and pinpoint how it was brought into the school, our staff is in the process of conducting a thorough inspection of the school and treating appropriately where necessary. We will continue to monitor the situation and consult with public health and pest control professionals to eliminate any bed bugs in the building and to minimize the potential for future bed bug activity in the school.

Although bed bugs are a nuisance, they have not been known to spread disease. Anyone who comes in direct contact with bed bugs or their eggs can unknowingly carry them into their home or workplace. They are easily spread by moving beds, furniture, luggage, or clothing from one location to another. For this reason, it is important that you inspect your child's backpack, lunch bag and other items regularly for the presence of bedbugs.

It is recommended that you routinely check your residence for signs of bed bug activity. Eliminating an extensive bed bug infestation can be a very challenging and costly process, so it is very important to begin treatments and prevention practices as soon as a problem is identified.

Bed bugs require blood to survive and reproduce, so they will typically look to bite any areas of exposed skin while you are asleep. The bites may not hurt at first, but may become swollen and itch a few days later, much like a mosquito bite. Most bed bug bites do not usually necessitate medical treatment. Wash the affected area with soap and water and then apply Calamine lotion to help dry out the bumps and stop the itching. If allergic reactions persist, an antibiotic cream or topical steroid, such as Hydrocortisone, will help take care of the infection. Please contact your physician with specific medical questions.

If you have any questions regarding bed bugs or our response efforts, please feel free to contact the Principal or School Nurse. Detailed information on bed bugs and appropriate control measures can be obtained on the Cuyahoga County Board of Health's web site (www.ccbh.net). Thank you for your assistance in this matter.

Sincerely,

Principal

BED BUGS: IDENTIFICATION, INSPECTION & TREATMENT



Public health officials and pest control operators in northeast Ohio and across the country are responding to a significant increase in bed bug complaints. Bed bugs were once found only on bedding in homes, apartments, and rooming houses. Now they are found in office buildings, retail stores, hospitals, dormitories, nursing homes, libraries, movie theaters, buses, or any other place where people gather. The Centers for Disease Control (CDC) and the Environmental Protection Agency (EPA) have declared that the bed bug is now a “pest of significant public health importance”.

Identifying Bed Bugs

- Bed bugs are small, flat, oval, reddish-brown, wingless insects that feed on the blood of humans and other mammals.
- Adult bed bugs are approximately ¼ inch long, about the size of an apple seed. Young bed bugs (nymphs) are quite small and when unfed they appear lighter and almost clear in color.
- Bed bugs do not fly or jump. However, they can crawl very fast.
- Female bed bugs can lay up to five eggs a day and five hundred during a lifetime.
- They are excellent hitchhikers and are easily spread by moving beds, furniture, luggage, clothes, or other possessions from one location to another.
- Anyone who comes in direct contact with bed bugs or their eggs can unknowingly carry them into their home or workplace.



Bed bug eggs and cast skins



Fecal spots by outlet cover

Bed Bug Bites

- Bed bugs are primarily a nuisance to humans and are not known to transmit disease.
- Bites can cause welts which may become infected by excessive scratching.
- Bed bug infestations can cause anxiety as well as psychological and emotional stress.



Welts from bed bug bites

Signs of a Bed Bug Infestation

Usually the first sign of a bed bug infestation is the appearance of red itchy welts on any bare skin that is exposed while sleeping. Be aware that not every welt or bite is due to bed bugs. Consult a medical provider with questions about unexplained bites or red welts.

Live bed bugs, eggs and cast skins indicate a bed bug infestation. Small black or rusty colored spots found on bed linens, pillows, or the mattress may be blood spots and bed bug droppings.



Blood stains and fecal spots on a mattress

Inspecting for Bed Bugs


Bed bugs hide close to where people sleep. They prefer fabric, wood, and paper surfaces over metal or plastic. Look for live bed bugs, eggs, cast skins, and blood or fecal spots in these locations:

- | | | | | |
|--|--|--|---|--|
| <input type="checkbox"/> mattresses | <input type="checkbox"/> recliners | <input type="checkbox"/> telephones | <input type="checkbox"/> luggage | <input type="checkbox"/> hollow furniture legs |
| <input type="checkbox"/> box springs | <input type="checkbox"/> baseboards | <input type="checkbox"/> radios | <input type="checkbox"/> futons | <input type="checkbox"/> door frames & hinges |
| <input type="checkbox"/> head boards | <input type="checkbox"/> behind pictures | <input type="checkbox"/> televisions | <input type="checkbox"/> gym bags | <input type="checkbox"/> wall / ceiling junction |
| <input type="checkbox"/> bed frames | <input type="checkbox"/> under loose wallpaper | <input type="checkbox"/> stacks of books | <input type="checkbox"/> draperies & curtains | <input type="checkbox"/> wheelchairs/walkers |
| <input type="checkbox"/> upholstered furniture | <input type="checkbox"/> sleeper sofas | <input type="checkbox"/> piles of papers | <input type="checkbox"/> stuffed animals | <input type="checkbox"/> bed side tables |
| | <input type="checkbox"/> electrical outlets | <input type="checkbox"/> back packs | | <input type="checkbox"/> edge of carpeting |

Treating Bed Bug Infestations

Complete elimination of a bed bug infestation can be a difficult process and may require the services of a knowledgeable and licensed pest management company. It may take several treatments to gain control over an infestation. If a “do-it-yourself” method is chosen, only use pesticide products that are labeled to kill bed bugs. Always read and follow the label directions before applying any pesticide product. Here are some additional tips to help eliminate bed bugs.

- Reduce and eliminate clutter. Don't keep piles of clothes, boxes, toys, shoes, etc. on the floor, under the bed, or in closets. They are prime hiding places for bed bugs.
- Heat kills bed bugs. Wash clothes in the hottest recommended setting. Tumble dry the clothes on high heat for at least 30 minutes (120° F or above).
- Encase an infested mattress and box spring in a zippered cover that is labeled and certified “bed bug proof”. Leave the covers on for at least one full year.
- Vacuum your home thoroughly and often. Pay particular attention to the area around the bed and the bed itself. Sprinkle ¼ to ½ cup of talcum powder on the last section of floor to be vacuumed. The powder will help dry out and kill bed bugs in the bag or cup. Place the vacuum cleaner bag or contents in a sealable plastic bag and discard it in the trash outside.
- Pesticides labeled to kill bed bugs are available over the counter and may provide effective control. However, if the problem persists or is heavily entrenched, contact a knowledgeable, experienced, and licensed pest management professional for assistance.
- Since bed bugs are difficult to control, plan on several thorough treatments to eliminate an infestation.
- DO NOT USE “Bug Bombs”. These products may kill on contact but they are mostly ineffective against hidden bed bugs. They may make the infestation worse by scattering the bugs throughout the home or apartment.
- If an infestation is suspected in a rental unit, contact the building manager or landlord about the problem. Property owners should contact a professional pest control company for advice and assistance. Your local health department can also provide assistance on bed bug identification, prevention and control options.
- The property manager and pest management professional may provide preparation guidelines prior to treating for bed bugs. Getting rid of bed bugs is a cooperative effort. And these guidelines should be followed closely.



Wash sheets and blankets in hot water and dry them thoroughly in a hot dryer before making up the bed



A zippered bed bug proof cover can help protect against bed bugs



Do not use foggers

Preventing Future Infestations of Bed Bugs

- Do not bring discarded bed frames, mattresses, box springs, or upholstered furniture into the home.
- Carefully inspect used or rented furniture prior to bringing it into the home.
- Place any recently purchased clothes, whether new or second hand, into the dryer on a hot setting for at least 30 minutes (120°F or above).
- When traveling, inspect the bed, headboard, and furniture upon arrival. Keep suitcases off the floor and bed and inspect them before leaving. Wash and dry all clothing, per the above recommendations, after returning home.
- Caulk and seal any cracks and crevices throughout the home, especially in rooms where people sleep.
- Be careful of who stays overnight or sleeps at the house. Likewise, be mindful when staying away from home. Inspect all children’s backpacks, stuffed animals, and other belongings if they have stayed overnight or spent considerable time at another location.





Extension FactSheet

Entomology, 1991 Kenny Road, Columbus, OH 43210

Bed Bugs

Susan C. Jones, Ph.D.,
Assistant Professor of Entomology
Extension Specialist, Household & Structural Pests

Bed bugs are parasites that preferentially feed on humans. If people aren't available, they instead will feed on other warm-blooded animals, including birds, rodents, bats, and pets.

Bed bugs have been documented as pests since the 17th century. They were introduced into our country by the early colonists. Bed bugs were common in the United States prior to World War II, after which time widespread use of synthetic insecticides such as DDT greatly reduced their numbers. Improvements in household and personal cleanliness as well as increased regulation of the used furniture market also likely contributed to their reduced pest status.

In the past decade, bed bugs have begun making a comeback across the United States, although they are not considered to be a major pest. The widespread use of baits rather than insecticide sprays for ant and cockroach control is a factor that has been implicated in their return. Bed bugs are blood feeders that do not feed on ant and cockroach baits. International travel and commerce are thought to facilitate the spread of these insect hitchhikers, because eggs, young, and adult bed bugs are readily transported in luggage, clothing, bedding, and furniture. Bed bugs can infest airplanes, ships, trains, and buses. Bed bugs are most frequently found in dwellings with a high rate of occupant turnover, such as hotels, motels, hostels, dormitories, shelters, apartment complexes, tenements, and prisons. Such infestations usually are not a reflection of poor hygiene or bad housekeeping.

Distribution

Bed bugs are fairly cosmopolitan. *Cimex lectularius* is most frequently found in the northern temperate climates of North America, Europe, and Central Asia, although it occurs sporadically in southern temperate regions. The tropical bed bug, *C. hemipterus*, is adapted for semitropical to tropical climates and is widespread in the warmer areas of Africa, Asia, and the tropics of North America and South America. In the United States, *C. hemipterus* occurs in Florida.



Mature Bed Bug

Order: Family—Hemiptera: Cimicidae	
Common name	Scientific name
Bed Bug	<i>Cimex lectularius</i>
Tropical Bed Bug	<i>Cimex hemipterus</i>

Identification

Adult bed bugs are brown to reddish-brown, oval-shaped, flattened, and about 3/16 to 1/5 inch long. Their flat shape enables them to readily hide in cracks and crevices. The body becomes more elongate, swollen, and dark red after a blood meal. Bed bugs have a beaklike piercing-sucking mouthpart system. The adults have small, stubby, nonfunctional wing pads. Newly hatched nymphs are nearly colorless, becoming brownish as they mature.

Nymphs have the general appearance of adults. Eggs are white and about 1/32 inch long.

Bed bugs superficially resemble a number of closely related insects (family Cimicidae), such as bat bugs (*Cimex adjunctus*), chimney swift bugs (*Cimexopsis* spp.), and swallow bugs (*Oeciacus* spp.). A microscope is needed to examine the insect for distinguishing characteristics, which often requires the skills of an entomologist. In Ohio, bat bugs are far more common than bed bugs.

Life Cycle

Female bed bugs lay from one to twelve eggs per day, and the eggs are deposited on rough surfaces or in crack and crevices. The eggs are coated with a sticky substance so they adhere to the substrate. Eggs hatch in 6 to 17 days, and nymphs can immediately begin to feed. They require a blood meal in order to molt. Bed bugs reach maturity after five molts. Developmental time (egg to adult) is affected by temperature and takes about 21 days at 86° F to 120 days at 65° F. The nymphal period is greatly prolonged when food is scarce. Nymphs and adults can live for several months without food. The adult's lifespan may encompass 12-18 months. Three or more generations can occur each year.

Habits

Bed bugs are fast moving insects that are nocturnal blood-feeders. They feed mostly at night when their host is asleep. After using their sharp beak to pierce the skin of a host, they inject a salivary fluid containing an anticoagulant that helps them obtain blood. Nymphs may become engorged with blood within three minutes, whereas a full-grown bed bug usually feeds for ten to fifteen minutes. They then crawl away to a hiding place to digest the meal. When hungry, bed bugs again search for a host.

Bed bugs hide during the day in dark, protected sites. They seem to prefer fabric, wood, and paper surfaces. They usually occur in fairly close proximity to the host, although they can travel far distances. Bed bugs initially can be found about tufts, seams, and folds of mattresses, later spreading to crevices in the bedstead. In heavier infestations, they also may occupy hiding places farther from the bed. They may hide in window and door frames, electrical boxes, floor cracks, baseboards, furniture, and under the tack board of wall-to-wall carpeting. Bed bugs often crawl upward to hide in pictures, wall hangings, drapery pleats, loosened wallpaper, cracks in plaster, and ceiling moldings.

Injury

The bite is painless. The salivary fluid injected by bed bugs typically causes the skin to become irritated and inflamed, although individuals can differ in their sensitivity. A small, hard, swollen, white welt may develop at the site of each bite. This is accompanied by severe itching that lasts for several hours to days. Scratching may cause the welts to become infected. The

amount of blood loss due to bed bug feeding typically does not adversely affect the host.

Rows of three or so welts on exposed skin are characteristic signs of bed bugs. Welts do not have a red spot in the center such as is characteristic of flea bites.

Some individuals respond to bed bug infestations with anxiety, stress, and insomnia. Bed bugs are not known to transmit disease.

Tell-tale Signs

A bed bug infestation can be recognized by blood stains from crushed bugs or by rusty (sometimes dark) spots of excrement on sheets and mattresses, bed clothes, and walls. Fecal spots, eggshells, and shed skins may be found in the vicinity of their hiding places. An offensive, sweet, musty odor from their scent glands may be detected when bed bug infestations are severe.

Control Measures

A critical first step is to correctly identify the blood-feeding pest, as this determines which management tactics to adopt that take into account specific bug biology and habits. For example, if the blood-feeder is a bat bug rather than a bed bug, a different management approach is needed.

Control of bed bugs is best achieved by following an integrated pest management (IPM) approach that involves multiple tactics, such as preventive measures, sanitation, and chemicals applied to targeted sites. Severe infestations usually are best handled by a licensed pest management professional.

Prevention

Do not bring infested items into one's home. It is important to carefully inspect clothing and baggage of travelers, being on the lookout for bed bugs and their tell-tale fecal spots. Also, inspect secondhand beds, bedding, and furniture. Caulk cracks and crevices in the building exterior and also repair or screen openings to exclude birds, bats, and rodents that can serve as alternate hosts for bed bugs.

Inspection

A thorough inspection of the premises to locate bed bugs and their harborage sites is necessary so that cleaning efforts and insecticide treatments can be focused. Inspection efforts should concentrate on the mattress, box springs, and bed frame, as well as crack and crevices that the bed bugs may hide in during the day or when digesting a blood meal. The latter sites include window and door frames, floor cracks, carpet tack boards, baseboards, electrical boxes, furniture, pictures, wall hangings, drapery pleats, loosened wallpaper, cracks in plaster, and ceiling moldings. Determine whether birds or rodents are nesting on or near the house.

In hotels, apartments, and other multiple-type dwellings, it

is advisable to also inspect adjoining units since bed bugs can travel long distances.

Sanitation

Sanitation measures include frequently vacuuming the mattress and premises, laundering bedding and clothing in hot water, and cleaning and sanitizing dwellings. After vacuuming, immediately place the vacuum cleaner bag in a plastic bag, seal tightly, and discard in a container outdoors—this prevents captured bed bugs from escaping into the home. A stiff brush can be used to scrub the mattress seams to dislodge bed bugs and eggs. Discarding the mattress is another option, although a new mattress can quickly become infested if bed bugs are still on the premises. Steam cleaning of mattresses generally is not recommended because it is difficult to get rid of excess moisture, which can lead to problems with mold, mildew, house dust mites, etc.

Repair cracks in plaster and glue down loosened wallpaper to eliminate bed bug harborage sites. Remove and destroy wild animal roosts and nests when possible.

Trapping

After the mattress is vacuumed or scrubbed, it can be enclosed in a zippered mattress cover such as that used for house dust mites. Any bed bugs remaining on the mattress will be trapped inside the cover. Leave the cover in place for a year or so since bed bugs can live for a long time without a blood meal.

Sticky traps or glueboards may be used to capture bed bugs that wander about. However, the effectiveness of these traps is

not well documented.

Insecticides

Residual insecticides (usually pyrethroids) are applied as spot treatments to cracks and crevices where bed bugs are hiding. Increased penetration of the insecticide into cracks and crevices can be achieved if accumulated dirt and debris are first removed using a vacuum cleaner. Avoid using highly repellent formulations, which cause bed bugs to scatter to many places. Dust formulations may be used to treat wall voids and attics. Repeat insecticide applications if bed bugs are present two weeks after the initial treatment since it is difficult to find all hiding places and hidden eggs may have hatched.

Do not use any insecticide on a mattress unless the product label specifically mentions such use. Note that very few insecticides are labeled for use on mattresses. If using an appropriately labeled insecticide on a mattress, take measures to minimize pesticide exposure to occupants. Apply the insecticide as a light mist to the entire mattress, opening seams, tufts, and folds to allow the chemical to penetrate into these hiding areas. Allow the treated surface to completely dry before use. Do not sleep directly on a treated mattress; be sure bed linens are in place. Do not treat mattresses of infants or ill people. Alternatives to using an insecticide on a mattress are discussed in the ‘Sanitation’ and ‘Trapping’ sections.

No insecticides are labeled for use on bedding or linens. These items should be dry cleaned or laundered in hot water and dried using the “hot” setting.

Visit Ohio State University Extension’s WWW site “Ohioline”
at:
<http://ohioline.osu.edu>

All educational programs conducted by Ohio State University Extension are available to clientele on a nondiscriminatory basis without regard to race, color, creed, religion, sexual orientation, national origin, gender, age, disability or Vietnam-era veteran status.

Keith L. Smith, Associate Vice President for Ag. Adm. and Director, OSU Extension

TDD No. 800-589-8292 (Ohio only) or 614-292-1868

1/2004-des

